

1940.

QUEENSLAND.

FORTIETH ANNUAL REPORT OF THE BUREAU  
OF SUGAR EXPERIMENT STATIONS.

REPORT OF THE DIRECTOR

TO

THE HON. THE SECRETARY FOR AGRICULTURE AND STOCK

(As required by "The Sugar Experiment Stations Acts, 1900 to 1938").

PRESENTED TO PARLIAMENT BY COMMAND.

BRISBANE:

BY AUTHORITY: A. H. TUCKER, GOVERNMENT PRINTER.

A. 36—1940.

n.t. Sugar.

1937.

6-373

6-889

6-834

7-010

6-902

6-765

6-416

6-593

6-709

6-902

6-755

6-426

6-256

6-319

6-511

6-481

6-537

6-620

6-584

6-698

6-340

6-418

6-415

6-443

7-877

7-710

7-128

7-826

8-357

7-574

7-536

7-850

7-608

8-052

11-585

7-603



# FORTIETH ANNUAL REPORT OF THE BUREAU OF SUGAR EXPERIMENT STATIONS.

## Director's Report.

TO THE HONOURABLE THE SECRETARY FOR AGRICULTURE AND STOCK.

SIR,—I have the honour to submit the Fortieth Annual Report of The Bureau of Sugar Experiment Stations, covering the period 1st July, 1939, to 30th June, 1940.

H. W. KERR,  
Director.

Brisbane, 1st November, 1940.

### GENERAL.

The growing season for the 1940 cane crop was one of alternating extremes. The dry spring months of late 1939 were followed, in general, by excessively heavy wet season rains, so that water-logging of fields of backward cane was common, and the crop was handicapped due to this cause. The magnitude of the estimated crop is therefore surprisingly good, having regard for the season.

In the far-Northern districts, very little rain fell during the latter half of 1939; at Meringa, this yielded a total of 3.63 inches. Many fields possessed a drought-stricken appearance when the wet season rains arrived. Canes on the higher lands benefited from the prolonged wet conditions, but all crops arrowed freely, in common with the experience of other districts of the State.

The Central districts experienced a somewhat similar season. In these parts, the wet season was short, and growth was checked after the last favourable rain period of early April.

In the Southern districts, the crop was produced under conditions very similar to those of the Central districts. A prolonged dry autumn was followed by damaging frosts in the winter.

The cyclone season also left its mark on the crop. The areas from Mackay to Cairns were affected, in part, by one or more cyclonic storms. The most severe damage occurred in the Burdekin district, where an abnormal flood, accompanied by wind, was responsible for much crop loss. The earlier crop estimates in these parts had therefore to be revised very considerably.

#### CROP YIELD, 1939—CROP ESTIMATE, 1940.

The following table shows the individual mill crushings for the 1938 season, and the estimated crops available for harvest during 1939:—

1939 Crushing.	Mill.	1940 Estimate.
Tons.		Tons.
130,063	Mossman .. .. .	137,000
212,011	Hambledon .. .. .	225,000
271,008	Mulgrave .. .. .	260,000
216,711	Babinda .. .. .	240,000
181,315	Goondi .. .. .	192,000
254,195	South Johnstone .. .. .	250,000
196,734	Mourilyan .. .. .	187,000
302,594	Tully .. .. .	280,000
319,297	Victoria .. .. .	240,000
295,071	Macknade .. .. .	240,000
124,200	Invicta .. .. .	105,000
198,556	Pioneer .. .. .	184,950
243,815	Kalamia .. .. .	180,000
272,024	Inkerman .. .. .	216,000
164,533	Proserpine .. .. .	160,000
96,733	Cattle Creek .. .. .	85,000
242,894	Racecourse .. .. .	215,000
218,942	Farleigh .. .. .	211,000
128,010	North Eton .. .. .	130,000
230,207	Marian .. .. .	185,000
245,958	Pleystowe .. .. .	218,500
231,396	Plane Creek .. .. .	218,000
91,127	Qunaba .. .. .	94,000
191,197	Millaquin .. .. .	240,000
224,075	Bingera .. .. .	280,000
231,981	Fairyhead .. .. .	290,000
60,623	Gin Gin .. .. .	71,150
211,836	Isis .. .. .	261,500
62,017	Maryborough .. .. .	65,000
50,379	Mount Bauple .. .. .	48,000
119,312	Moreton .. .. .	149,500
17,711	Rocky Point .. .. .	16,500
2,296	Eagleby .. .. .	2,600
6,038,821.	Total .. .. .	5,857,700

#### ESTIMATES OF SUGAR YIELD, 1940 CROP.

The preliminary estimate of the cane crop which will be available for harvest in Queensland during the 1940 season is 5,857,700 tons. This is substantially below the actual yield for 1939, but is, nevertheless, very much higher than normal. At the present juncture it is doubtful whether this tonnage will be realised, due to the heavy arrowing of the crop in most districts, combined with lighter crop yields following the very dry autumn. On the other hand, the crop will probably show a high sugar content, and this factor will compensate in some measure.

Allowing for that proportion of the crop in the Southern areas which will probably be allowed to stand over until 1941, it is anticipated that some 810,000 tons of raw sugar will be manufactured. Such a tonnage, if realised, will be second in magnitude only to the record production of 1939—891,422 tons.

In addition, it is estimated that the New South Wales mills will crush some 310,000 tons of cane, from which approximately 39,000 tons of sugar may be expected. The estimated total Australian production for 1940 may therefore be placed at 850,000 tons.

#### STATISTICS OF THE 1939 CROP.

The yield of raw sugar in Queensland for the 1939 crop was 891,422 tons\* of 94 n.t. This was easily an all-time record tonnage, exceeding that of 1938 by some 113,000 tons.

The following table shows the geographical distribution of the crop for the past five years, as between "northern" and "southern" cane districts:—

*Sugar Production, 1935-'39.*

District.	1935.	1936.	1937.	1938.	1939.
North of Townsville	Tons. 258,958	Tons. 333,615	Tons. 373,692	Tons. 328,301	Tons. 351,267
South of Townsville	.. 351,368	.. 410,646	.. 389,633	.. 449,835	.. 540,155
Total .. .. .	610,326	744,261	763,325	778,136	891,422

The progressive upward trend in production in the central and southern areas of the State has thus been maintained, and over 60 per cent. of the record tonnage was produced from areas south of Townsville.

These results are due in part to each of the following causes:—(1) Planting of superior cane varieties, (2) elimination of loss due to diseases, (3) increased acres cultivated, and (4) the adoption of better methods of cultivation and more rational use of manures.

#### ACRES HARVESTED AND ACREAGE YIELD.

The total area harvested for milling purposes in 1939 was 261,047 acres, which is almost exactly 10,000 acres greater than that of the previous season. The acreages under plant, ratoon, and standover crops were as follows:—

Plant cane .. .. .	Acres. 102,759
Ratoon cane .. .. .	.. 137,590
Standover cane .. .. .	.. 20,698
Total .. .. .	261,047

\* This is slightly in excess of the figure quoted on page 2, as it includes "local" sales which are additional to that acquired by the Sugar Board.

The yield of cane per acre crushed was 23.1 tons, while the average sugar yield was 3.41 tons. Both these figures establish new records for Queensland, and the yield of sugar per acre has now exceeded 3 tons for four consecutive years.

The following were the yields of cane and sugar per acre in the respective sugar districts, during 1939:—

District.	Tons Cane per acre.	Tons 94 n.t. Sugar per Acre.
Mossman-Ingham .. .. .	23.2	3.43
Lower Burdekin .. .. .	29.5	4.71
Proserpine .. .. .	15.0	2.36
Mackay-St. Lawrence .. .. .	19.7	2.99
Bundaberg-Gin Gin .. .. .	28.0	3.71
Maryborough-Childers-Gympie .. .. .	22.6	3.10
Nambour-Beenleigh .. .. .	25.3	3.34
State Average .. .. .	23.14	3.41

The following table gives the crop statistics for Queensland for the past ten years:—

Table showing Acres Cultivated and Harvested, Yields of Cane and Sugar, Acre-Yields, and Quality of Cane, 1930-1939.

Year.	Acres Cultivated.	Acres Harvested.	Total Yields.		Yields per Acre.		Tons Cane to 1 Ton Sugar.
			Cane.	Sugar.	Cane.	Sugar.	
1930 ..	296,070	222,044	Tons. 3,528,660	Tons. 516,783	Tons. 15.89	Tons. 2.33	6.83
1931 ..	309,818	233,304	4,034,300	581,276	17.29	2.49	6.94
1932 ..	291,136	205,046	3,546,443	514,085	17.30	2.51	6.90
1933 ..	311,910	228,154	4,667,028	638,734	20.46	2.80	7.31
1934 ..	303,916	218,126	4,269,991	612,570	19.56	2.80	6.97
1935 ..	314,700	228,515	4,220,267	610,326	18.47	2.67	6.92
1936 ..	338,686	245,152	5,171,516	744,261	21.10	3.04	6.94
1937 ..	348,840	249,683	5,132,934	763,325	20.56	3.06	6.73
1938 ..	347,199	251,064	5,342,085	778,136	21.28	3.10	6.87
1939 ..	*	261,047	6,038,821	891,422	23.14	3.41	6.77
True Average for 10 Years ..	234,244	4,595,205	665,092	19.62	2.84	6.91	

\* Not available.

AVERAGE AREA HARVESTED PER FARM.

The following figures show the average acreage harvested by cane planters in Queensland for the 1938 crop:—

	Acres.
Cairns to Townsville .. .. .	48
Ayr to Mackay .. .. .	35
Bundaberg to Bauple .. .. .	20
Nambour to Beenleigh .. .. .	8
State average .. .. .	32

The average area harvested per planter was 32 acres, which is one acre higher than that of the previous year.

MOLASSES PRODUCED.

The following figures supplied by the Government Statistician show the manner in which the molasses produced in 1939 was disposed of:—

	Gallons.
Sold to distilleries .. .. .	9,581,241
Burnt as fuel .. .. .	3,834,653
Used or sold for feed .. .. .	4,727,170
Sold or used for other purposes .. .. .	188,889
Used as manure .. .. .	4,295,289
Run to waste .. .. .	457,541
Total .. .. .	23,084,783

ECONOMIC REVIEW.

The 1939 crushing season was launched in an atmosphere of uncertainty. The cane crop available for harvest was estimated at more than 6 million tons—the largest ever produced. With the operation of the international export quota scheme, it appeared that less than 50,000 tons of excess sugar could be acquired and disposed of, over and above the basic mill peak aggregate of 737,000 tons. The probable unmarketable excess was thus some 100,000 tons of raw sugar.

At this period the grave international situation was rapidly deteriorating, and culminated in the declaration of war early in September. This brought about a substantial change in the economic situation notably insofar as the British Empire was concerned. Besides being released, as belligerent countries, from the control of the International Sugar Agreement, the British Government embarked on a policy of Empire purchasing, which ultimately gave the Australian sugar producer the unexpected opportunity for disposing of his entire 1939 production. The transaction was, of course, contingent upon the availability of shipping for the transportation of raw sugar to those part of the British Empire where it was required.

The benefits of such a liberalised marketing offer was fully availed of by all producers from Mackay north; the Southern Queensland growers, though harvesting in excess of normal limits, still elected to pursue their policy of standing over portion of the excess cane until the following year. The

MAFFRA BEET FACTORY.

The following data in respect of the Maffra Sugar Factory, Victoria, are supplied through the courtesy of the Manager:—

Area harvested .. .. .	3,950 acres
Beet purchased .. .. .	42,898 tons
Beet sliced .. .. .	41,890 tons
Average sugar content .. .. .	18.86 per cent.
Sugar produced .. .. .	6,280 tons
Price paid for beet .. .. .	46/-
Average yield beet per acre .. .. .	10.86 tons
Average yield refined sugar per acre .. .. .	1.59 tons

1939 SUGAR VALUES.

The proportion of the 1939 sugar crop manufactured in Queensland, which was required for consumption and use in the Commonwealth of Australia, was declared at 49.8495 per cent., and for that export at 50.1505 per cent. These proportions are exclusive of the "excess" sugar produced by mills in excess of their allotments under the Peak Year Scheme.† The excess sugar produced for the 1939 season was 168,520 tons, as compared with 163,943 tons for the 1938 crop.

The price payable for the sugar required for consumption and use in Australia was declared at £23 12s. 6d. per ton of 94 net titre. The net value per ton of 94 net titre sugar sold abroad was £10 7s. 6d., which is £2 3s. 3d. per ton higher than the 1938 figure. The average price paid to those Queensland mills which did not produce "excess" sugar was £16 19s. 7d. per ton, compared with £16 19s. 1d. for the previous season.‡ The average value of all sugar was £15 15s. 3d., which is the highest recorded since 1935.

The following table summarises production and consumption figures and sugar values since the year 1924, when the first large surplus was produced:—

Year.	Total Sugar Production at 94 n.t.	Tons Sugar Exported.*	Average Australian Price.	Average Export Price.	Average Price, No. 1 Pool Sugar.	Average Price, all Sugar.
1924 ..	Tons. 409,136	Tons. 74,000	£ 26.0	£ 21.0	£ 26.0	£ 26.0
1925 ..	485,585	219,000	26.5	11.3	19.5	19.5
1926 ..	389,272	74,777	26.5	14.9	24.5	24.5
1927 ..	485,745	152,384	26.5	12.1	22.0	22.0
1928 ..	520,620	186,703	26.5	10.5	20.9	20.9
1929 ..	518,516	197,000	27.0	9.9	20.3	20.3
1930 ..	516,783	203,605	27.0	8.3	19.7	19.5
1931 ..	581,276	291,802	27.0	9.4	18.3	18.0
1932 ..	514,027	189,733	25.0	8.3	19.3	18.8
1933 ..	638,734	305,687	24.0	8.0	17.2	16.2
1934 ..	612,570	277,336	24.0	7.6	16.5	15.5
1935 ..	610,326	298,202	24.0	7.9	16.9	16.2
1936 ..	744,261	409,400	24.1	7.95	17.1	15.2
1937 ..	762,794	430,523	24.0	8.3	17.55	15.3
1938 ..	776,810	443,386	24.0	8.2	16.95	15.1
1939 ..	890,896	515,792	23.6	10.4	17.0	15.75

\* Bagged sugar. † Peak Year Scheme first operated in 1930.

‡ Revised Mill Peak schedule first operated in 1939.

TOTAL VALUE, 1939 SUGAR CROP.

The total value of the 1939 Queensland crop was £14,042,000—an all-time record, and higher than the 1938 value by £2,304,000.

net export value realised for sugar sold after the outbreak of hostilities was about £10 10s. per ton, which is substantially better than the average price of recent years.

The aggregate sugar tonnage manufactured was 891,000 valued at over £14,000,000. Both figures constitute record high values for Queensland sugar production.

While the war was thus initially responsible for easing what would doubtless have been a most difficult position, there are definite indications that the continuance of hostilities may create just as difficult problems for the future. The early estimates of the 1940 cane crop suggest that the tonnage which will be available for manufacture is capable of yielding well over 800,000 tons of sugar, and the reduction on the 1939 harvest is due only to the less favourable growing season experienced. While it has been announced that the Empire purchasing plan may be able to absorb the entire volume of production during 1940, the availability of shipping is causing no little concern. It was anticipated that the sugar manufactured early in the crushing season could be got away without difficulty, but it appeared probable that a large proportion of that manufactured later in the season would have to be held in storage for some months. Many of the mills were forced to store portion of the 1939 crop until the autumn of 1940, and steps have since been taken to provide increased storage capacity for this and subsequent years. Doubtless the British Government will utilize all available ships

in such a ma commodities in appear that st served, and th of this commo

The indu the need for acute than ex sincere effort of the Regu provision for on individual tonnage. Th majority of of some nu scheme while react to the

Howeve thing has l mill areas to the indi a net acrea possesses tl It mus control is

The Divisions in the es summari of the F

Me were he Sth Ma the Dej March

St in an : positio applica decide Techn that t nation

I staff were Elliot Messr not t office

I the F is at

Boa the oper Boa suec area has stat pro gro bee tio: sue

wh un gr at ha st

fe

in such a manner as to grant preferential rights to those commodities in most urgent demand. In this respect it would appear that stocks of sugar in England have been well conserved, and there is no suggestion of any immediate shortage of this commodity.

The industry recognises that the outlook is obscure, and the need for rationalisation of production is probably more acute than ever before. In most mill areas of the State a sincere effort has been made to implement the 1939 Amendment of the Regulation of Sugar Cane Prices Acts, which makes provision for the formulation of a scheme to control production on individual farms within the limits of the present mill peak tonnage. To devise a basis which is acceptable to the large majority of canegrowers in any area is, of course, a task of some magnitude. This is inevitable, for an alternative scheme which favours one group of farmers must automatically react to the detriment of the rest.

However, it is encouraging to be able to record that something has been achieved in this respect. While a number of mill areas have allotted peak tonnages of cane (or sugar) to the individual suppliers, others have favoured control on a net acreage basis. The future alone will show which plan possesses the greater merit.

It must be admitted frankly that any form of production control is irksome to the primary producer. Even though

he may have no special desire to increase his individual production, there exists the factor of intensification of agricultural methods which makes for a steady but progressive increase in acreage yield. While it may be urged that the area cultivated should be reduced in proportion to the increased production per acre, this would be offset by the natural gain in regional farming population from year to year, which normally prefers to be absorbed into sugar production rather than migrate to an area where other forms of primary industry are enjoying no greater measure of stability.

Considerations such as these have induced certain of the leaders of the industry to explore the possibilities of utilizing any such surplus production for purposes other than human consumption, as well as providing a safety valve in the event of the closing of the present overseas distribution channels which are at present available, during the period of post-war adjustment. The scheme of diverting cane juices or syrups for the purpose of fuel alcohol production seems to offer something in this regard.

Admittedly alcohol produced from farm crops cannot compete economically with imported fuel, so long as present supplies of petrol are available; but perhaps the manufacture of a proportion of the Nation's liquid fuel requirements from home-grown produce will provide scope for rationalisation in some fields of primary production which are at present facing a most difficult problem.

WORK OF THE BUREAU.

The detailed reports of the officers in charge of the several Divisions and of regional Experiment Stations will be found in the ensuing pages of this Report. The following comments summarise the more important general aspects of the activities of the Bureau for the year under review:—

ADVISORY BOARD.

*Meetings.*—During the year, two meetings of the Board were held in Brisbane—on the 14th December, 1939, and on the 8th March, 1940. In the absence of the Hon. the Minister, the Deputy Chairman (Mr. N. H. Wellard) presided over the March meeting.

*Staff.*—During the year further applications were called in an attempt to fill the vacancy of Mill Technologist. The position was advertised throughout the sugar world, and applications were received from several countries. The Board decided to appoint Mr. E. R. Behne, Senior Assistant Mill Technologist, to the position. It was recommended, further, that this officer be sent abroad for a period, when the international position has returned to normal.

In pursuance of the Board's policy to build up the field staff to a satisfactory working strength, further two cadets were appointed to the staff in the persons of Messrs. J. T. Elliott and E. V. Humphry. Unfortunately, the services of Messrs. E. J. Barke, G. A. Christie, and E. V. Humphry will not be available for a period, due to the enlistment of these officers with the fighting forces.

Mr. L. C. Home, who has served as a part-time officer of the Bureau, has now been appointed Laboratory Assistant, and is attached to the Brisbane chemical laboratory.

*Disease and Pest Control.*—Much of the time of the Board's meetings was occupied in considering matters affecting the more adequate control of pests and diseases. Most of the operations of the recently constituted Cane Disease Control Boards have fully justified the hopes which were held for the success of this project as an aid to disease control. In some areas it may justly be claimed that the vigilance of the boards has been responsible for a very marked improvement in the status of disease control, and in at least one area, it is probable that the major variety would have been lost to the growers if the control measures initiated by the Board had not been taken. These boards generally function in close co-operation with the Bureau; this is, of course, essential to their success and smooth working.

*Non-approved Cane Varieties.*—Despite the wide publicity which was given to cane-growers respecting their obligations under the Amending Act of October, 1938, and the year of grace which extended to all farmers who had planted non-approved canes, it was found that a large number of growers had failed to take adequate steps for the destruction of such fields or odd stools of canes.

To deal with these recalcitrants has imposed upon the field staff of the Bureau a particularly onerous task, the

difficulties of which were magnified due to loss of staff through enlistment. The Board is very definitely agreed that the requirements of the Act shall be observed by all growers, if it is to be effective in its operation.

*Amendment of Act.*—Following representations by canegrowers' organisations, the Board agreed that where pests and disease control boards were operating in any area, they should be merged into one with full control over both phases. It is anticipated that the Act will be amended during the forthcoming Parliamentary session to take care of this. A further proposal deals with the period during which varieties "approved when planted but subsequently disapproved" may be cultivated, to meet the peculiar circumstances operating in certain districts.

*Field Days.*—Farmers' Field Days were conducted on each of the three Experiment Stations. The Mackay field day was held in April, in conjunction with the meetings of the Queensland Society of Sugar Cane Technologists, while the Meringa and Bundaberg functions were held in June. In each case the attendances were satisfactory.

*New Cane Varieties.*—The Board again endorsed the policy of the Bureau in seeking to improve the variety status of the industry, through the breeding and importation of new canes. During the year, certain new varieties bred in Queensland were released for trial purposes, while others are undergoing further observational tests prior to their being propagated on a commercial scale.

*Publications.*—During the year the "Queensland Cane Growers' Handbook" was issued, and a bound copy of the publication was forwarded to each canegrower in the State. This has been favourably received and commented upon by farmers generally. The "Cane Growers' Quarterly Bulletin" and "Technical Communications" have been continued, while a periodical "News Letter" is now being issued for the guidance of the mills and their technical staffs.

*Utilization of By-products.*—The Board devoted further attention to this subject, notably in reference to the attempts which are being made to utilize molasses as a substitute for bitumen in road construction. The efforts which are being made to expand the power alcohol industry in this State are also receiving the careful scrutiny of the Board.

*Levy.*—The levy for the year 1940-41 was fixed at 4d. per ton of cane, as in the previous year.

QUEENSLAND SOCIETY OF SUGAR CANE TECHNOLOGISTS.

The 1940 Conference of the Society was held in Mackay. The meetings were a complete success in every particular, and it would be difficult to over-estimate the value of this sister organisation in its contributions to the well-being of the industry. It is strongly supported by the technical officers of the Bureau, who contributed their quota of papers as in former years.

R.Y.  
Sugarcane Sugar Factory,  
of the Manager:—

950 acres  
898 tons  
890 tons  
-86 per cent.  
280 tons  
/-  
86 tons  
59 tons

manufactured in  
tion and use in  
d at 49,8495 per  
cent. These pro-  
gar produced by  
the Peak Year  
1939 season was  
or the 1938 crop.

for consumption  
2s. 6d. per ton  
net titre sugar  
. per ton higher  
o those Queens-  
s" sugar was  
s. 1d. for the  
all sugar was  
ice 1935.

and consump-  
1924, when the

Average Price, No. 1 Pool Sugar.	Average Price, all Sugar.
£ 26.0	£ 26.0
19.5	19.5
24.5	24.5
22.0	22.0
20.9	20.9
20.3	20.3
19.7	19.5
18.3	18.0
19.3	18.8
17.2	16.2
6.5	15.5
6.9	16.2
7.1	15.2
-55	15.3
-95	15.1
7.0	15.75

n 1930.  
crop was  
the 1938

break of  
stantially

s 891,000  
cord high

r easing  
position,  
ostilities  
e. The  
tonnage  
yielding  
on the  
r season  
Empire  
lume of  
causing  
anufac-  
without  
portion  
to be  
s were  
return  
reased  
btless  
ships

BALANCE-SHEET.

STATEMENT OF RECEIPTS AND DISBURSEMENTS FROM 1ST JULY, 1939, TO 30TH JUNE, 1940.

RECEIPTS.			DISBURSEMENTS.		
	£	s. d.		£	s. d.
To Balance, 1st July, 1939	18,114	15 6	By Salaries	10,734	15 2
" Assessments	18,875	17 6	" Contingencies	9,250	4 0
" Endowment	7,000	0 0	" Bundaberg Contingencies	1,806	11 9
" Bundaberg Stations	1,138	10 0	" Mackay Contingencies	1,782	18 11
" Mackay Station	1,029	18 2	" Meringa Contingencies	2,098	13 4
" Meringa Station	634	7 11	" Balance, 30th June, 1940	21,502	17 10
" Sundries	382	11 11			
	<u>£47,176</u>	<u>1 0</u>		<u>£47,176</u>	<u>1 0</u>

STATEMENT OF RECEIPTS AND DISBURSEMENTS FROM INCEPTION OF FUND, 1ST DECEMBER, 1900, TO 30TH JUNE, 1940.

RECEIPTS.			DISBURSEMENTS.		
	£	s. d.		£	s. d.
To Assessments	268,958	7 4	By Disbursements	507,179	5 11
" Endowment	206,199	1 7	" Balance	21,502	17 10
" Sugar Experiment Stations	53,524	14 10			
	<u>£528,682</u>	<u>3 9</u>		<u>£528,682</u>	<u>3 9</u>

C.C.S. FORMULA.

For the benefit of overseas readers of this report, we would point out that the measure of available sugar in cane (so-called Commercial Cane Sugar or C.C.S.) is defined as follows:—

$$\text{C.C.S. in juice} = \frac{\text{Pol in juice} - \frac{\text{Impurities in juice}}{2}}{2}$$

and C.C.S. in cane is calculated by the formula:—

$$\text{C.C.S.} = \frac{3P}{2} \left(1 - \frac{5 + F}{100}\right) - \frac{B}{2} \left(1 - \frac{3 + F}{100}\right)$$

where—P = pol in first expressed juice  
 B = brix in first expressed juice  
 F = fibre in cane.

DIVISION OF SOILS AND AGRICULTURE.

WORK OF THE BRISBANE LABORATORY.

MR. C. R. VON STIEGLITZ, Chemist.

*Routine Analyses.*—The following is a summary of the routine analyses performed at the Brisbane Laboratory for the period 1st July, 1939, to 30th June, 1940:—

Soils	547
Waters	29
Limes	11
Sugar Canes	11
Sugars	5
Molasses	2
Fertilizers	2
Miscellaneous	3
<b>Total</b>	<b>610</b>

The number of soil samples submitted by farmers for fertilizer advice, based on analytical results, increased 32 per cent. over the previous year's total, but farmers have not availed themselves of this free service to the extent anticipated. The addition of Mr. L. C. Home to the Bureau staff during the year has provided much needed help in handling more expeditiously soil samples forwarded for fertilizer advice, and has enabled more time to be utilized in special analytical studies and investigational work generally.

*Laboratory Training of Field Officers.*—Messrs. Smith, Elliott, and Humphry, Cadets, spent varying periods of instruction in the Brisbane Laboratory spread over approximately three months. In addition to gaining experience in the simpler analytical methods used for examining soils for plant food deficiencies, experiments were designed to exemplify the different physical, chemical, and biological characteristics of major soil types. These experiments were carried out by the Cadets themselves under the supervision of one of the chemical staff.

*Investigational Work.*—"Technical Communication" No. 10, 1939, "Basic Lead Acetate: Its Composition and Clarifying Qualities," by C. R. von Stieglitz, was published during the year. This publication lists analyses of twenty-two samples of basic lead acetate as used at Queensland sugar mills, and describes experiments employed for assessing the clarifying efficiency of such.

*Soil Fertility Survey.*—Work was continued on No. 6 Branch, South Johnstone area, in collaboration with Mr. Knust, instructor in cane culture, and the results will be mapped as previously. The further sampling of this area is to be discontinued, however, as the widespread application of mill by-products to these soils has rendered difficult a correct interpretation of the analytical results.

Investigations designed to furnish information on possible loss of soil nutrients through leaching, with particular reference to potash, were commenced early in the year. Soils from

different districts are being sampled monthly and analysed for available potash and phosphate, and pH. Three soil types have been selected from the Immisfail district, one from Mackay, and one from Bundaberg.

LABORATORY METHODS OF ASSESSING AVAILABLE PLANT.

*Nutrients.*—The usual yearly comparison between the results of farm fertility trials and laboratory methods for determining available phosphate and potash was carried out, and resulted in fourteen out of seventeen, or 82 per cent., of the cases being correctly placed for both plant foods. Of these three exceptions, one soil showed high acidity which might well have been the limiting factor, and of the other two one was distinctly a border line case for phosphate. The results may therefore be regarded as highly satisfactory and confirm previous findings.

*Nitrogen Availability.*—Investigational work with a view to finding a suitable method of assessing the availability of this plant food in soils was continued, and a modification of Andrews' method of estimating the carbon dioxide produced by soil organisms, when certain optimum conditions are observed, shows promise of providing useful data. The method, however, would not seem to have the general application of the phosphate and potash tests, but when the soils are grouped according to districts and major soil types an examination of the available data suggests that a useful correlation will be established.

*Damage to Soils by Cyclone.*—The flooding of fields in certain areas of the Burdekin Delta, due to cyclonic weather in the early months of the year, resulted in extensive damage to crops and fields. Attention has been drawn to the extent of this damage by Dr. H. W. Kerr in the July issue of the "Queensland Cane Growers' Quarterly Bulletin." As a result of his inspection of the district, a number of soil samples was forwarded to Brisbane for chemical investigation; some represented the subsoils of fields which had suffered extensive erosion and others silt or sand which had been deposited on the original soil.

The result of the chemical analyses demonstrated that reserves of the soil plant foods, phosphate and potash, of these soils are quite satisfactory in the majority of cases, even in the sandy deposits, but the total nitrogen and nitrate producing power, except in the case of two silt deposits, are very low. The chief problems facing the farmer in these affected areas will be the regrading of the fields to permit irrigation practices to operate successfully and the most economic utilization of such irrigation where the sandy deposits are superimposed on the original surface soils.

METEORO

However seasons may accentuated total rainfa closely app amount fell remainder f uneven dist effect on g were almos period othe however, a that were growth th that were retarded ir

As a : tracted we during th satisfactor experiee considerat absence o of the soi Later, wi showed co when ligh of overca tion; her falls wer were ins and the was extr crops an their ap quickly rapid ar

AF

July, 19  
 August,  
 Septemb  
 October,  
 Novemb  
 Decemb  
 January  
 Februar  
 March,  
 April, 1  
 May, 19  
 June, 19

1  
 canes  
 2  
 again

again

G.13  
 G.1:  
 G.1:  
 G.1  
 G.2  
 H.2  
 S.J

NORTHERN SUGAR EXPERIMENT STATION, MERINGA.

Mr. R. W. MUNGOMERY, Acting Officer in Charge.

METEOROLOGICAL, AND NOTES ON CROP GROWTH.

However marked the contrast between the wet and dry seasons may have been during 1938-39, it was even more accentuated during the 1939-40 growing period. Although a total rainfall of 82.10 inches was recorded (and this very closely approximates the average) only 3.63 inches of this amount fell during the last six months of 1939, whilst the remainder fell on 107 wet days in the first half of 1940. This uneven distribution of rainfall could not have but a detrimental effect on growth; for in the spring months some of the fields were almost drought stricken, whereas during the autumn period other fields were waterlogged. Taken by and large, however, a crop only slightly below average resulted, for crops that were backward on the higher lands made satisfactory growth throughout the continued rainy season, whilst those that were forward on the lower ground were somewhat retarded in growth as the soil became saturated with water.

As a result of the copious rains that fell during the protracted wet season of 1939, soil moistures generally were good during the late winter months, and conditions were quite satisfactory for late plantings; but very cold weather was experienced in August, and in consequence, germinations were considerably delayed. In addition, high winds and the absence of any further useful rains soon caused a depletion of the soil moisture, and the cane was slow in coming away. Later, with the persistence of dry conditions, the young cane showed considerable distress until towards the end of October, when light showers fell. These were accompanied by a period of overcast weather, and little moisture was lost by evaporation; hence the young cane made a rapid recovery. Scanty falls were received at intervals during November, but these were insufficient to allow of vigorous growth taking place, and the absence of substantial rain during December, which was extremely hot, caused a severe check in the growth of all crops and they wilted badly. However, thunderstorms made their appearance early in January, and soil moistures were quickly replenished, so that from then onwards recovery was rapid and a vigorous growth was maintained.

On 18th February, this Experiment Station came under the influence of a cyclonic disturbance, and winds of gale force blew down trees and completely flattened some of the more advanced crops. Other damage occurred through some of the more vigorously growing varieties snapping off at the succulent internode just below the top, and this later caused considerable side shooting. Fortunately the disturbance was accompanied by very little rain, and the crops were generally not in a forward state, otherwise the damage would have been much greater. Monsoonal rains and showery conditions continued right into June; as a result it was not practicable to cultivate some of the land required for the early plantings, and much of the anticipated early plantings had to be deferred. In addition, a considerable proportion of the cane was still immature when harvesting operations commenced. A quantity of grub-damaged cane from neighbouring districts was received at the mill, but grub damage did not occur on this station. Prolific arrowing occurred during May, and some of the backward and late cut crops were seriously affected.

The following are the rainfall records taken at this Experiment Station during the past twenty years:—

Year.	Rainfall in Inches.	Year.	Rainfall in Inches.
1920	94.86	1931	98.82
1921	122.84	1932	76.31
1922	64.90	1933	95.06
1923	53.29	1934	91.44
1924	95.67	1935	59.91
1925	76.98	1936	88.81
1926	59.12	1937	46.33
1927	90.16	1938	55.86
1928	66.33	1939	118.08
1929	102.28	1940 (6 months)	79.47
1930	107.61	Average, 24 years	81.66

ABSTRACT OF METEOROLOGICAL OBSERVATIONS MADE AT THE NORTHERN SUGAR EXPERIMENT STATION, MERINGA, FROM 1ST JULY, 1939, TO 30TH JUNE, 1940.

Month.	Rainfall in Inches.	Number of Wet Days.	Highest Shade Maximum.	Lowest Shade Maximum.	Mean Shade Maximum.	Highest Shade Minimum.	Lowest Shade Minimum.	Mean Shade Minimum.	Mean Diurnal Range.	Mean Temperature 9 a.m.	Mean Relative Humidity 9 a.m.
July, 1939	.32	5	85.0	72.4	78.3	67.3	43.0	57.2	21.2	67.9	84
August, 1939	.28	2	90.4	74.0	81.0	64.6	37.4	48.7	32.3	67.3	78
September, 1939	.25	6	92.0	78.8	84.3	64.2	42.4	56.3	28.0	73.2	77
October, 1939	.90	7	93.0	78.7	87.3	71.6	45.6	62.9	24.3	78.2	68
November, 1939	1.63	7	95.6	80.0	90.6	75.6	63.4	69.3	21.2	82.2	69
December, 1939	.25	2	99.2	90.2	94.6	74.5	63.3	70.0	24.7	85.9	59
January, 1940	12.67	14	101.9	78.0	91.6	77.4	68.7	72.8	18.7	82.6	80
February, 1940	18.87	21	96.7	84.2	90.9	77.5	71.1	74.9	16.0	82.0	80
March, 1940	34.37	25	96.1	80.3	96.7	75.5	64.2	72.0	14.7	79.2	87
April, 1940	7.27	17	96.0	80.2	85.2	75.0	58.9	69.1	16.2	78.4	82
May, 1940	1.70	11	87.2	79.1	83.1	69.5	46.8	64.2	18.8	74.6	81
June, 1940	3.59	19	84.0	71.0	78.8	69.0	49.0	62.9	15.8	72.5	84
Totals	82.10	136	..	..	..	..	..	..	..	..	..

NEW EXPERIMENTS INITIATED DURING THE YEAR.

1. Varietal Trial.—5 × 5 Latin square, 4 new seedling canes v. S.J.4.
2. Observational Yield Trial.—Sixteen advanced seedlings against a standard variety.

EXPERIMENTS HARVESTED DURING 1939.

1. Observational Yield Trial.—Six advanced seedlings against S.J.4.

OBSERVATIONAL VARIETAL TRIAL.

Varieties.—G.113, 118, 126, 140, 243, H.248, and S.J.4.  
 Area of Plot.—0.05 acre.  
 Harvested.—November, 1939.  
 Age of Crop.—15½ months.  
 Crop Yields.—(Individual plots of seedling canes.)

Variety.	Cane per Acre.	C.C.S. in Cane.
G.113	Tons.	Per cent.
G.118	27.4	15.0*
G.126	37.4	14.5
G.140	31.4	17.5
G.243	38.0	16.8
H.248	24.4	18.6
S.J.4 (3 plots)	28.6	15.5
	32.6	17.0

\* From maturity tests.

DISCUSSION.

Four of these seedlings—G.113, G.126, G.243, and H.248—were selected earlier in the year and planted in a Latin square trial in July, 1940. The results of this trial will be available during the coming crushing season. On tonnage figures it would appear that G.140 and G.118 should have been selected; however, the former showed high susceptibility to leaf scald disease, and was discarded on that account, and G.118 was eliminated on bad habit and very low sugar content during most of the period up to harvest.

The observational trial was ratooned and fertilized, and the first ratoon crop yields will be obtained during the 1940 season.

LAND CLEARING AND NEW PLANTINGS.

A further area of forest land was cleared during the year, in order to provide space for new experiments, chiefly associated with seedling propagation and trials. Some two acres of new land were brought under the plough, while work was done on existing blocks to provide better drainage conditions.

With the promise attending preliminary attempts to transfer cross pollination work to Meringa from Freshwater, more extensive varietal plantings have been made on the station to provide the parental canes for future seasons.

INTRODUCTION OF NEW VARIETIES.

Several new varieties were planted on the Station in 1939. These comprised promising seedlings raised in other

£	s.	d.
10,734	15	2
9,250	4	0
1,806	11	9
1,782	18	11
2,098	13	4
21,502	17	10
£47,176	1	0

1940.

£	s.	d.
507,179	5	11
21,502	17	10
£528,682	3	9

sugar in cane

analysed for oil types have from Mackay,

E PLANT. Between the methods for carried out, per cent, of foods. Of idity which f the other phate. The factory and

with a view lability of fication of produced itions are ie method, ion of the grouped nation of n will be

fields in weather damage ie extent e of the a result ples was ie repre- xtensive sited on

ed that ash, of s, even nitrate its, are these permit most deposits

States or by the Colonial Sugar Refining Co. Ltd. in their nurseries, as well as canes from other parts of the world which had just been released from the Brisbane quarantine house. They included:—

C.P.807	31-1389	30 S.N.1031
C.P.29/116	30 R.115	33 S.N.1160
P.O.J.2875	30 G.1250	37 N.G.6
B.726	30 G.1759	Atlas
B.2935	30 S.N.225	Cato
Q.20	30 S.N.362	Comus
Q.25	30 S.N.451	Marcus
Q.29	30 S.N.874	

In all cases, only a few setts of each variety were available, and the crop therefrom will be utilized in planting disease-resistance trials as well as further propagation plots.

LEGUMES.

The search for valuable leguminous species suitable for these parts was continued. Through the courtesy of the Council for Scientific and Industrial Research and other organisations, a wide range of new species was planted during the year. Those under trial included the following:—*Crotalaria maxillaris*, *usaramoensis*, *anagyroides*, *juncea*, *Stizolobium atterinum*, *pachylobium*, *Dolichos lab-lab*, *debilis*, *Phaseolus riccardianus* (strains C.P.I.2778 and 2779), *trinervis*, *Sesbania paulensis*, *Glycine javanica*, *Centrosema pubescens*, *Calapogonium mucunoides*, and *Pueraria javanica*.

Plantings made in the spring were handicapped by dry weather, but they made good headway following the summer thunderstorms. The summer plantings received good rains at the outset, and made continued growth until late in the season.

*Crotalaria usaramoensis* gave an excellent germination and seems to possess a better early coverage than other species of the same genus under trial here, whilst it has a good long growing period. Hence it shows more than ordinary promise, and more extensive plantings will be made during the coming year.

*Crotalaria anagyroides* made good growth and gave a heavy crop of green matter when ploughed under, though it tends to become rather woody as it ages. It likewise appears to be promising.

*Phaseolus riccardianus* (2 strains) and *P. trinervis* all germinated well, and, despite the relatively dry conditions during their early growth period, they grew vigorously and produced a good volume of green matter before commencing to seed in the autumn.

*Pueraria javanica* vined and spread well, and produced an abundant crop of green matter. These plants did not flower and commence to set seed until late May, hence this legume should be of special value to those growers who require the protection of a green crop on their lands until well into the winter months.

Further confirmatory plantings of the above legumes will be made during the coming year, and, in addition, plantings will be made of *Stizolobium atterinum*, *S. pachylobium*, and *Dolichos lab-lab*, which although they produced fair crops, were somewhat affected by the prolonged wet conditions since the plots of these legumes happened to be located on ground somewhat lower than the remainder. The other species will be discarded.

FODDER CROPS.

Due to the risk attaching to plantings of maize in cane areas where downy mildew exists, experiments are being conducted at this Station with a view to determining the fodder value of alternative species. A number of grain sorghums were tested for this purpose during the past year, and preliminary results are very encouraging. Varieties Schrock, Coleman, Atlas, American Early Red, Hegari, and Manko were included in the trial; this list also places the varieties approximately in their order of merit. Schrock was definitely outstanding in all three plantings. They produced good heads of grain, and remained free of peach moth caterpillars which are reported to be troublesome in other areas of the State.

Brazilian lucerne (*Stilosanthes guyannensis*) was also introduced and propagated. It produced particularly good

yields on some of the poorer forest country, and it should prove a valuable complementary species when sown on some of the poor land grass paddocks.

COTTON PLANTINGS.

In co-operation with the Cotton Branch of the Department of Agriculture, a small trial planting of cotton was made on this Station, to determine the possibilities of the crop in the humid tropics. The varieties included in the trial were—Half and Half, Miller, Okra Acala, Oklahoma Triumph, and New Brooklyn. Planting was done on the 23rd August, 1939.

Owing to the dry spring, germinations were erratic and growth very backward until November. It was not until the end of December that the first bolls opened, and the majority opened during the wet season; there was, therefore, considerable waste of cotton. Insect damage was slight, but stainers become plentiful later in the season.

From these observations it would not appear that cotton-growing will be found a reliable venture in the wet coastal belt, unless the crop can be forced to maturity either before or after the wet season rains. If before, it is likely that earlier planting combined with irrigation would be necessary.

LABORATORY WORK.

In addition to the customary routine work of sampling new varieties both on the Station and in farm trials, soil analyses for fertilizer and lime requirements were undertaken at this laboratory during the last year, and in this connection a fairly intensive soil sampling programme was carried out in the Mossman area, on those farms where the growers were requiring this service. The following samples were analysed during the year:—

	Samples.
Canes (Station analyses and general maturity tests)	719
Canes, farm trials	118
Canes, farmers' samples	11
Soils	102
<b>Totals</b>	<b>950</b>

FIELD DAY.

The second annual Field Day was held on this station on 8th June, 1940, and although the attendance was not up to the standard that might be expected from an area of this size, still it was 50 per cent. greater than that of the previous year; this in itself was encouraging. However, several who did attend came from distant centres, such as Innisfail and Mossman, and all growers exhibited keenness to keep *au fait* with the progress in this branch of their industry. They were welcomed by the Director and the Deputy Chairman of the Advisory Board, after which followed addresses by the Director and members of his staff on topical agricultural problems. A tour of inspection of the station was then made, and growers generally were interested in viewing the new variety introductions, the locally bred seedlings, the new legumes, and the crops of grain sorghums, which were in excellent head at this particular time.

CROP SUMMARY.

	Tons.
Cane sent to the mill	471
Cane used for plants, &c.	13
<b>Total</b>	<b>484</b>
	<b>Acres.</b>
Total area harvested	14.2
Tons cane per acre harvested	34.1

An analysis of the yields shows the following allocation in respect of classes and varieties:—

	Tons.
Plant cane	296
Ratoon cane	188
<b>Total</b>	<b>484</b>
<b>Varieties—</b>	
Badilla	104
S.J.4	131
Q.10	87
Miscellaneous (including seedlings)	162
<b>Total</b>	<b>484</b>

The rainfall in excess of that during February, 1940, on the yield for the year for the waterlogged, and

The period for dry. December a start, and prosper

ABSTRACT OF ME

Month.

	1939.	1940
July	..	..
August	..	..
September	..	..
October	..	..
November	..	..
December	..	..
January	..	..
February	..	..
March	..	..
April	..	..
May	..	..
June	..	..

ANNUAL RAINF.

Year

1920	..
1921	..
1922	..
1923	..
1924	..
1925	..
1926	..
1927	..
1928	..
1929	..
1930	..

Ex

1. Va

S

2. Ob

C

3. Dc

1. V:

2. V

3. C

4. I



CENTRAL SUGAR EXPERIMENT STATION, MACKAY.

Mr. D. L. MCBRYDE, Chemist in Charge.

METEOROLOGICAL.

The rainfall for the 1939-40 growing season was much in excess of that of the previous year...

The period from July to November, 1939, was exceedingly dry. December and January rains gave the young crop a good start...

bright until the wet season proper set in. The season was unusual, in that five cyclonic disturbances affected the district.

The early winter was mild and dry, and crops made little progress during the period. Crop estimates for 1940 are therefore substantially below the record harvest...

ABSTRACT OF METEOROLOGICAL OBSERVATIONS MADE AT THE SUGAR EXPERIMENT STATION AT MACKAY, FROM 1ST JULY, 1939, TO 30TH JUNE, 1940.

Table with columns for Month, Inches Rainfall, Wet Days, Average Rainfall, and Shade Temperatures (Maximum, Minimum, High, Low, Mean).

N.B.—Rainfalls—(a) = average 39 years, (b) = average 40 years.

ANNUAL RAINFALL SINCE 1920 AT SUGAR EXPERIMENT STATION, MACKAY.

Table showing annual rainfall in inches from 1920 to 1939, with average values for 39 and 40 years.

Fertilizer Applied.—All plots received 4 cwt. Sugar Bureau No. 1 Planting Mixture in the drill at the time of planting...

SUMMARY OF YIELDS, PLANT CROP.

Table with columns for Variety, Cane per Acre (Tons), and C.C.S. in Cane (Per cent).

DISCUSSION.

Varieties E.4, E.45, and Q.28 are three seedlings which were raised at the Mackay Station and which survived final selection.

EXPERIMENTS HARVESTED DURING 1939 SEASON.

- 1. Varietal Trial—Mackay Seedlings v. Q.813, Latin Square, Plant Cane.
2. Observational Variety Trial—Mackay Seedlings v. Q.813, Single Plot, Plant Cane.
3. Downy Mildew Resistance Trial.

EXPERIMENTS INITIATED DURING 1939.

- 1. Varietal Trial—Mackay Seedlings v. Q813, Latin Square, Plant Cane.
2. Varietal Trial—Jason, Comus, Q.20, Q.28 v. Q.813, Latin Square, Plant Cane.
3. Observational Variety Trial—Seedlings v. Q.813, Plant Cane.
4. Fertility Trial—Factorial Trial (3 x 3 x 3), Plant Cane.

VARIETAL TRIAL (Plant Crop).

PLAN AND YIELDS.

Table showing yields for varieties E.45, Q.813, Comus, Q.28, and E.4.

OBSERVATIONAL VARIETAL TRIAL (Plant Crop).

PLAN AND YIELDS.

Table showing yields for varieties F.57, F.21, Q.813, G.22, G.39, G.18, G.19, F.40, G.5, G.52, Q.813, Q.813, G.58, F.58, G.17, G.20, G.8, Q.813, F.31, and Q.813.

Block.—B2.

Harvested.—October, 1939.

Age of Crop.—14 months.

Experimental Plan.—Single plots with contacting Q.813 plots.

Plots.—1/20 acre.

Fertilizer Applied.—4 cwt. Sugar Bureau No. 1 Planting Mixture in the drill, and 3 cwt. of sulphate of ammonia in two top dressings...

Block.—B2.

Harvested.—October, 1939.

Age of Crop.—14 months.

Experimental Plan.—5 x 5 Latin Square.

Plots.—1/18 acre.





TREATMENT AND GROWTH NOTES.

This observational yield trial was planted in August, 1938, on land which had a 12 months' fallow under peas and partly under Gambia pea. The block is not a very good one owing to ironstone gravel near the surface. The crop was backward from the start, and even at time of harvesting did not appear to be carrying the crop which the weights indicate. It was fertilized with Sugar Bureau Planting Mixture No. 3 at the rate of 3 cwt. per acre.

CROP YIELDS.

Table with columns: Variety, Cane per Acre., C.C.S. in Cane. Rows include F.28, F.25, F.26, F.10, Co.290 (3 plots), F.19, F.22, F.23, F.21, F.20.

DISCUSSION.

This preliminary trial embracing advanced seedlings was set out for the purpose of enabling the more promising to be selected for inclusion in a Latin square yield trial. As individual plots only were possible, the potential value of the varieties must be assessed on other qualities as well as yield index. Several apparently outyielded the standard (Co.290). In all cases sugar values were rather low; most varieties were quite immature when harvested, but this was unavoidable as the plots had to provide planting material for a further trial.

FACTORIAL FERTILITY TRIAL (First Ratoon Crop). PLAN AND YIELDS.

Factorial Fertility Trial table showing combinations of treatments (1N, 1P, 0K, 2N, 2P, 2K, 0N, 1N, 2N, etc.) and their corresponding yields.

Block.—B1. Harvested.—October, 1939. Age of Crop.—13 months. Variety.—Co.290. Plan.—3 x 3 x 3 factorial trial. Plots.—0.007 acre.

TREATMENT AND GROWTH NOTES.

Subsequent to harvesting the plant crop the block was ratooned, as usual, with the single-tyne subsoiler—three times per interspace. Fertilizer was then applied at the following rates:—

- 1N = 120 lb. sulphate of ammonia per acre.
2N = 240 lb. sulphate of ammonia per acre.
1P = 400 lb. superphosphate per acre.
2P = 800 lb. superphosphate per acre.
1K = 180 lb. muriate of potash per acre.
2K = 360 lb. muriate of potash per acre.

All the phosphate and potash was applied at ratooning, while the sulphate of ammonia was applied in two equal dressings when the crop was further advanced, and weather conditions suitable. The crop ratooned well, but was badly wilted and wind-burnt during December and January. It revived with late January rains, and proceeded to make good growth in the later months.

SUMMARY OF YIELDS—PLANT AND FIRST RATOON CROPS.

Summary of Yields table comparing Plant Crop and First Ratoon Crop across different treatments (0N, 1N, 2N, 0P, 1P, 2P, 0K, 1K, 2K).

DISCUSSION.

As was the case with the plant crop, the first ratoon yields exhibited no benefits from any of the plant-food treatments. The influence of treatment on the C.C.S. values was, however, marked. With progressive potash applications, the sugar content of the crop was lowered, suggesting that the crop was over-mature when harvested. The heavy phosphate applications showed the opposite effect; where superphosphate was applied at the rate of 800 lb. per acre, the C.C.S. of the crop was the best. It is well recognised that one influence of phosphate on cane growth is to delay maturity.

This trial was again ratooned, and it is anticipated that the second ratoon yields will be very interesting.

VARIETAL TRIAL—Co.290 v. P.O.J.213 (First Ratoon Crop). PLAN AND YIELDS.

Varietal Trial table comparing yields of Co.290 and P.O.J.213 under various fertilizer treatments.

Block.—B2. Harvested.—October, 1939. Age of Crop.—14 months. Plots.—0.099 acre. Experimental Plan.—Nine randomised blocks.

TREATMENT.

This crop followed the harvesting of the plant crop in August, 1938. It was ratooned with the single-tyne subsoiler, and then fertilized with Sugar Bureau No. 3 ratooning mixture (4 cwt. per acre), followed later with 275 lb. of sulphate of ammonia per acre. During the dry summer months the P.O.J.213 established its lead, the Co.290 being severely wind-burnt and almost dead. Later in the year the lead of the P.O.J.213 did not seem so apparent, but it is evident in the yields.

SUMMARY OF

Table listing Varieties: Co.290, P.O.J.213.

The comparison is to be somewhat different; Co.290 over P.O.J.213 that in a dry in an average

VARI

Vertical list of variety and treatment combinations with yields: P.O.J.2878 (27.1), P.O.J.2875 (36.3), P.O.J.2725 (26.2), P.O.J.234 (23.3), P.O.J.2881 (31.4), P.O.J.294 (32.2), Co.290 (30.2).

Block. Harvested. Age of Crop. System. Plots.

This age of the harvesting usual with —and the mixture; a suitable acre.

From dry summer With the varieties time elapses variety had had lack of insofar climatic pathologies

V.

Vertical list of variety and treatment combinations with yields: P.O.J.28, P.O.J.2, P.O.J.28, Co.290, P.O.J.2, P.O.J.2, P.O.J.2.













## REPORT OF THE DIVISION OF ENTOMOLOGY AND PATHOLOGY.

Mr. ARTHUR F. BELL, Assistant Director.

### INTRODUCTION.

Last year we recorded the enactment of "The Sugar Experiment Stations Acts, 1900 to 1938," which introduced new and far-reaching amendments in the principles of legislative control of diseases and insect pests of sugar cane. Certain difficulties have naturally been encountered in the transition period from the old to the new regime, but conditions are rapidly becoming stabilised and the current season will, it is expected, see the transition completed. The benefits of more strict control of cane varieties have become obvious, while the ever-increasing yields of sugar per acre indicate that no detrimental economic effects result.

Following the upward trend of the past few years a favourable season resulted in a further increase in cane grub pest populations in North Queensland; damage was greater than it has been for some time past and, in the absence of soil fumigation campaigns, would have been very extensive. In view of the wartime difficulties of obtaining adequate fumigant supplies it is to be hoped that a climatic check will be imposed upon this pest this season.

The disease situation has continued to improve in Southern Queensland, chiefly owing to the expanded activities of Cane Disease Control Boards. These Boards have more than justified their constitution under the Sugar Experiment Stations Act, and are providing a valuable service to the sugar industry.

### CANE PESTS AND CANE DISEASE CONTROL BOARDS.

Two further Cane Disease Control Boards, viz., Mossman and Hambleton, were constituted during the year. Eight such Boards are now functioning; the activities are being directed in each case by qualified supervisors, who are bringing energy and enthusiasm to bear upon their work.

The sixth annual conference of Cane Pests Boards was held in Ayr on 12th June, 1940, and was attended by the record number of forty-four delegates, representing sixteen Boards. Nine papers dealing with various aspects of pest control were read and discussed. A considerable amount of interest centred upon an address by Mr. H. E. Lever, Technical Officer of Messrs. A. C. F. and Shirleys Limited, who was present, by invitation, to discuss the problem of obtaining adequate supplies of fumigants and poisons during wartime.

### THE GREYBACK CANE BEETLE (*Lepidoderma albobirtum* Waterh.).

The rapid increase of the greyback beetle in most of the Northern cane areas during the last year greatly outstripped the previous steady rate in the build-up of populations which was commented on in last year's report, and the grub position during the past season was such that serious losses occurred in many mill areas.

These losses did not occur despite fumigation, but rather through failure to fumigate. Although most of the mill areas have fallen into line and adopted the policy of soil fumigation as their method of front-line attack on the grub pest wherever serious infestations occur, some found their infested areas so great that available supplies of fumigant were totally inadequate to deal effectively with all the areas that needed attention, and supplies had to be rationed. Reserves that had been built up in other areas were freely drawn on until stocks were depleted, and the remaining untreated infestations were left to take their chance. Many fields therefore became almost a total loss, whilst others, in sections that received late deliveries of fumigant and were not treated at the optimum time, suffered serious checks. Another aspect which cannot be overlooked (and which, incidentally, will have an important bearing on next year's infestation) was the existence of extensive areas of light infestation which were not sufficiently severe to warrant fumigation but where, nevertheless, appreciable crop losses did occur. It has been estimated that the total acreage fumigated in North Queensland from Mossman to Tully was in the vicinity of 1,900 acres, which represents a considerable increase over that of previous years.

Fumigant deliveries were somewhat disorganised as a result of the war. This difficulty is expected to be overcome before the commencement of the next campaign by a co-operative effort on the part of the Pests Boards and the manufacturers, whereby orders will be placed early and supplies will be despatched whenever shipping space becomes available. The full quota for each mill area is therefore likely to be built up well in advance.

Considerable difficulty was experienced in the actual conduct of fumigation operations, due to the protracted beetle flights. This state of affairs is always likely to eventuate when early summer rains occur, and it places on the Supervisor the responsibility of advising the optimum time for fumigating mixed age groups.

Where the largest infestations resulted from early beetle flights, they were easily and effectively fumigated before the onset of the continuous monsoonal rains; but where the infestations were late the sodden condition of the soil rendered fumigation impossible, and necessitated delay until drier weather conditions were experienced, with consequent reduced efficiency. As a result of the development of the Blundell knapsack injector (referred to in a previous report) a large number of these units were used in various centres throughout the last fumigation campaign and the results were most gratifying. Not only did these injectors have less stoppages through mechanical defects or obstructions, but the dosages remained more constant; they were easier to manipulate, and the man-day performance was increased by approximately 100 per cent. over that obtained with the previous type. This serves to bring the campaign within limits closer to the optimum period or, alternatively, a larger acreage can be treated in the time previously occupied. Labour costs were reduced from approximately £2 10s. per acre to about £1 5s., and, as a result, these knapsack injectors are quickly displacing the older type of injector.

Speeding-up of the fumigating season has been carried a step further in the Mourilyan three-unit carriage machine by the provision of a battery-lighting set which flood-lights the base of the stools as the machine is drawn slowly along the cane rows, and so allows work to be undertaken during the night period, when conditions are cooler and more comfortable for the operators. The Sugar Workers' Award has been amended so that such work may proceed until midnight, making it possible to work two shifts with the one machine.

It is gratifying to note that some of those who were the greatest opponents of fumigation, when faced with ruin in the shape of ever-decreasing tonnages, low sucrose yields, and the replanting of large acreages annually, and having successfully tried fumigation, have now become some of the greatest advocates of this form of control. However, there still remain a few who appear loath to do anything to circumvent the pest other than by indirect methods, and it cannot be too strongly emphasised at this juncture that the large populations built up over extensive areas during the past year will, in their turn, give rise to even heavier infestations during the coming year if weather conditions should again prove favourable. Hence growers who have previously experienced depredations of this pest would be wise to heed this warning and make appropriate plans to cope with any anticipated infestation.

Attention has again been focussed on the cane area on the eastern side of the Basilisk Range where a tree-clearing project was initiated some few years ago, and it was of interest to note that many of the farms previously infested have, since the clearing of this range, remained free of the pest. At the same time farms to the south, where feeding trees still remain, and other farms elsewhere in the Mourilyan area have suffered heavy infestation. Such a state of affairs gives increasing support to the idea that this area will probably remain permanently free of the pest as a result of the elimination of the feeding trees.

Since this scheme has up to the present apparently justified itself, other tree-clearing schemes have been commenced, whilst others are being fostered or are under consideration. If successful, and not unduly costly, such tree-clearing projects eliminate for all time the necessity for undertaking any artificial control measures, but each scheme must be judged on its own merits. Undoubtedly areas of feeding trees from which canelands now become infested do occur, and by a co-operative effort some of these lands would profit by a destruction of the feeding trees. On the other hand, on account of the wide distribution of the feeding trees in some areas any scheme for the protection of the few adjacent canelands must surely prove uneconomic, and it must still be remembered that one of the earliest of such schemes proved a failure. All factors such as the possible fighting range of the beetles and the existence of any natural barriers, such as hills to obstruct their flight, should be taken into consideration when assessing the possible value of a projected scheme.

The elimination of the grub-resistant variety S.J.4 as a result of the outbreak of gumming disease in the Mulgrave and Hambleton districts has resulted in a concentrated search for replacement varieties. Trial plots of such varieties were examined to check their resistance to grub attack during the past season. Under the conditions of moderate infestations P.O.J.2725 exhibited a considerable degree of resistance; P.O.J.2878 showed less resistance than P.O.J.2725 but considerably more than the susceptible varieties Badila, Korpi, and Oramboo. Q.10 is susceptible and appears to attract larger numbers of beetles by virtue of its vigorous early growth habit; Q.29 on the other hand appears unattractive to beetles owing to its rather sparse early growth. Co.290

behaved similarly were made during a to be located near place near the sur reaction may be so

Canes which grubs suffer deterioration observed that the of the damage and tion of this factor and subjected to t simulate partial t March onwards; t Maturity tests co intervals. The C.C.S. for the spa canes; it would a in interpreting th

Infestations lighter than in t extended wet se reduced the num economic damage infestations of considerably more of heavy pest t Mount Jukes dis disappearance of were confined to dealt with by fu

On the othe of areas of gru of the very sus obvious damage the varieties it conclude that n Mackay farmers in blocks which to conduct annu

The Macka collecting of be season's main f then initiated i newly-appointed capacity by offic fumigated on districts. Adv fumigant, and t added to the d theless, the re number of far pating in this f of carbondisul the standard f being usually i

A consider the problem o attack to repl Mackay distric for grub resist cial varieties i

Resistan  
Interme  
Suscept.  
Ko

THE BEF

In most c depredations l been rather m that very we where the da incidence of t season as it l is probably tl troublesome. panies flood-normal level.

In acco horer-cum-tof Johnstone a Meringa Exp of top rot w not indicate any of the

From c operating in since it has which is dar than undam: that early m by severe g tends to ma

behaved similarly to P.O.J. 2878. The above observations were made during a particularly wet season when grubs tended to be located near the surface and thus root pruning took place near the surface; under drier conditions the relative reaction may be somewhat different.

Canes which have been subjected to root pruning by grubs suffer deterioration in juice quality but it has been observed that the rate of deterioration varies with the extent of the damage and from variety to variety. In an investigation of this factor fifteen varieties were planted at Meringa and subjected to three treatments, viz.: *a* check; *b* spaded (to simulate partial root pruning) at bi-weekly intervals from March onwards; *c* as in *b* but completely uprooted in May. Maturity tests commenced in May are being continued at intervals. The early analyses have shown an increase in C.C.S. for the spaded sections but a decrease for the uprooted canes; it would appear, however, that care must be exercised in interpreting the results.

Infestations in the Mackay district were generally much lighter than in the period 1936-39; on the other hand, the extended wet season, followed by continued dry weather, reduced the number of grubs per stool necessary to inflict economic damage. As a result, fields which experienced infestations of similar intensity in both years suffered considerably more damage this year. After a number of years of heavy pest populations and extensive grub damage the Mount Jukes district experienced a sudden and spectacular disappearance of general grub infestations; the few present were confined to relatively small areas and were adequately dealt with by fumigation.

On the other hand there were some apparent extensions of areas of grub damage due to recent extended plantings of the very susceptible variety S.J.2. This variety suffers obvious damage from populations which did not visually affect the varieties it has displaced and hence the tendency to conclude that new areas of grub infestation have occurred. Mackay farmers who contemplate further plantings of S.J.2 in blocks which might be infested by grubs should be prepared to conduct annual fumigation campaigns if necessary.

The Mackay Cane Pests Board ceased paying for the collecting of beetles early in January, 1940, and hence last season's main flights were not interfered with. The Board then initiated its first large-scale fumigation campaign, the newly-appointed Supervisor being assisted in an advisory capacity by officers of the Bureau. A total of 160 acres were fumigated on twenty-four farms distributed over several districts. Adverse weather conditions, irregular supplies of fumigant, and the occasion of more than one flight of beetles added to the difficulties of organising the campaign; nevertheless, the results were satisfactory and already a large number of farmers have signified their intention of participating in this form of grub control next season. The mixture of carbondisulphide and paradichlorobenzene has been made the standard fumigant in this district on account of there being usually more than one flight of beetles.

A considerable amount of attention has been devoted to the problem of obtaining varieties more resistant to grub attack to replace the discarded P.O.J.2878 in parts of the Mackay district and all new varieties are now being tested for grub resistance. The grub resistance of present commercial varieties in the Mackay district is as follows:—

*Resistant*: P.O.J.2878 and P.O.J.2714.

*Intermediate*: E.K.28.

*Susceptible*: 1900 Seedling, H.Q.426, Badila, Oramboo, Korpi, Q.813, Q.20, and S.J.2.

#### THE BEETLE BORER (*Rhabdocnemis obscura* Bois.).

In most of those areas that normally suffer badly from depredations by the beetle borer, damage from this pest has been rather moderate during the past year, in spite of the fact that very wet conditions prevailed. However, in localities where the damage from borers is generally most severe, the incidence of top rot was not nearly so marked during the past season as it has been in some of the previous years, and this is probably the reason why the borer pest did not prove more troublesome. Nevertheless the infestation that usually accompanies flood-damaged and rat-damaged cane remained at a normal level.

In accordance with the customary practice, a beetle borer-cum-top rot resistance trial was established in the South Johnstone area with third year seedling selections from Meringa Experiment Station; however, only a low percentage of top rot was recorded there and preharvest inspections did not indicate that there would be a high borer population in any of the varieties, including Badila.

From one locality reports were received of borers operating in grub-damaged cane. This was not unexpected since it has been observed on previous occasions that cane which is damaged by grubs is more susceptible to borer attack than undamaged cane. One possible reason for this is the fact that early maturity or over maturity and deterioration induced by severe grub attack exerts an attraction on the borers and tends to make them concentrate in the grub-damaged cane.

#### THE GIANT TOAD (*Bufo marinus* L.).

It would appear that populations of the Giant Toad have now reached saturation point in many Northern areas and although some growers affirm that at least in localised areas the toad is coping with the greyback beetle, in general it appears evident both on the score of observations in the field, and by virtue of the fact that grub damage has increased this year in areas where the toad population is high, that the number of beetles destroyed by the toad is relatively small when compared with the numbers that are nightly on the wing.

Examinations of the contents of toads' stomachs during the beetle fighting season were continued, and in a number of instances the remains of greyback beetles were found. However, even if the diet had consisted exclusively of beetles the number of toads present in the field was insufficient to effect a material reduction of the pest. This scarcity of toads in the field is not to be wondered at considering the absence of any debris or other cover which might serve as shelter for them during the day.

Evidence seems to indicate that the toad may be of more value in the reduction of beetle borer (*Rhabdocnemis obscura* Bois.). In the wetter areas where this pest thrives the toad population is more numerous in the canefields and it is believed probable that it may be particularly effective in dealing with migrating beetle borers.

#### THE SUGAR-CANE SCALE (*Aulacaspis madiunensis* Zehnt.).

Following reports of further severe infestations of the sugar-cane scale, *Aulacaspis madiunensis*, in the Bundaberg district, a comprehensive survey of the incidence and habits of this pest was undertaken in September, 1939. Information gleaned during this survey definitely establishes the existence of this scale in parts of the Bundaberg district prior to 1918. Fortunately it does not seem to have dispersed far from the older centres of infestation. Damage occurred in standover crops exclusively, and the insect was found to colonise only sparsely on crops up to the age of one year. One-year-old crops were standard practice up to some five or six years ago and it is considered that this explains why the pest made no headway and remained undetected. However, since the advent of irrigation, the success of P.O.J.2878 as a standover variety, and the attendant problems of over-production, standover cane has constituted a large percentage of the crop, and conditions have become favourable for the spread of the pest. Incidentally this demonstrates how the status of a minor pest can suddenly become altered under the conditions of a new economy.

The variety P.O.J.213 was the worst affected, followed by P.O.J.2878, whilst P.O.J.2725 and Co.290 appear to be fairly resistant to attack. Susceptibility or resistance to attack seems to follow no distinct breeding lines. Hence it was decided to put all promising new canes in South Queensland into a resistance trial to test their reaction to this pest, and a trial was accordingly established at South Kolan during the last spring.

Recommendations included the burning of affected blocks and the elimination of the variety P.O.J.213, which appears to have been largely instrumental in building up infestations. As this variety is more suited to a one-year cropping programme, and was fast losing the limited popularity it possessed, its loss will present no great hardships in the areas affected.

#### WIREWORMS (*Laeon variabilis* Cand.).

During the spring of 1939 many bad, indifferent, or slow strikes were experienced in canefields on several soil types in the Mackay district. Numerous field enquiries revealed that these were chiefly due to various combinations of unfavourable soil and climatic conditions and the varietal factor. Authentic wireworm infestations were relatively few and seldom severe and, consequently, a project for the study of field populations by mechanical soil sifting did not yield much useful information.

Over the period December, 1939-January, 1940, 52.23 inches of rain were recorded on forty-six wet days at the Mackay Experiment Station. A warning was therefore issued to farmers that severe and widespread damage could be expected during the forthcoming plantings. Actually, however, the autumn climatic conditions have been such that virtually no autumn planting has been carried out on wireworm country.

#### RATS.

During the past year the incidence of the rat pest in Queensland cane fields was again very low, and few fields showed evidence of appreciable economic damage. Most of the damage caused was due to the two species *Melomys littoralis* and *M. cervinipes*. Routine investigations of the problem of control were continued throughout the year. Some phases of this preliminary investigation have now been carried as far as is practicable and preparation of data for publication is in hand.





that the wet season was particularly heavy in Mackay this year, and normally dry areas approximated to the wet, swampy conditions characteristic of the section where dwarf disease has heretofore been found.

Red stripe and rind disease were less in evidence than usual in North Queensland, although at Mackay there was rather more red stripe than usual.

An experiment with the cross inoculation of possible strains of mosaic failed to give any evidence of the existence of any such differential strains.

#### CULTURES OF NITROGEN FIXING BACTERIA.

Some seventy-six canegrowers were supplied with Rhizobium cultures sufficient to inoculate 381 bushels of Poona pea seed, and 83 bushels of other cowpeas. Small quantities of culture were also supplied for the inoculation of lucerne, lupins, field peas, and Pueraria.

#### INTRODUCTION OF VARIETIES.

Importations of varieties from abroad comprised:—Uba Marot and Hind's Special from the United States; these canes will be used for breeding purposes following a recommendation by Mr. N. J. King on his return from the Louisiana Conference of the I.S.S.C.T. Varieties were also received from the Hawaiian Sugar Planters' Association as under:—

- 28-4291 (Hawaiian Uba x H.456).
- 31-2484 (P.O.J.2878 x (Uba x H.456)).
- 31-2806 (P.O.J.2878 x H.9811).
- 32-1063 (P.O.J.2878 x 28-4399 (= Hawaiian Uba x H.456)).
- 32-3575 (28-4898 (= Natal Uba x H.456) x 26.C.270).
- 32-8560 (Co.213 x P.O.J.2878).

The following varieties were also received from the Colonial Sugar Refining Company:—28R.154, 28R.155, 29G.706, 32G.1374, and 33S.N.1270.

## DIVISION OF MILL TECHNOLOGY.

Mr. E. R. BEHNE, Mill Technologist.

#### STAFF.

The classification of Mr. E. R. Behne was changed from that of Chief Assistant Technologist to that of Mill Technologist, whilst Mr. L. C. Home, who previously was Temporary Laboratory Assistant, was placed permanently on the staff.

#### MUTUAL CONTROL.

The Eighth Annual Synopsis of Mill Data for Mills in the Mutual Control, giving the figures for the 1939 season, has been published. Again twenty-four mills were incorporated in the scheme. The calculation sheets for the 1940 scheme have been revised and issued. The sheets are materially the same as those for the previous year, save for a minor alteration to the latter half of the reference numbers.

#### STANDARDISATION OF APPARATUS.

The following is a record of the standardisation work carried out during the year; this work forming an important part of the interseason activities:—

- Brix Spindles.*—473 spindles were tested, and all conformed to official requirements.
- Polariscope Tubes.*—33 were tested, 3 of which were unsatisfactory.
- Polariscopes.*—4 instruments were checked and adjusted.
- Pipettes.*—Of 191 pipettes tested, 30 were rejected.
- Burettes.*—13 were tested and found satisfactory.
- Cylinders.*—34 were tested and found satisfactory.
- Flasks.*—291 were tested, and 14 of these were condemned.
- Thermometers.*—3 were tested, all being satisfactory.
- Weights.*—6 sets were tested and adjusted when necessary.

In all, 1,048 pieces of apparatus were tested, compared with 578 for the previous year.

#### TECHNICAL PAPERS.

Six papers were presented by the Technology Staff of the Bureau at the Eleventh Annual Conference of the Queensland Society of Sugar Cane Technologists. These were as follows:—

- Behne, E. R., "The Commercial Cane Sugar (C.C.S.) Formula in Relation to Factory Control."
- Behne, E. R., "The Storage of Raw Sugar."
- Jenkins, G. H., "Cane Preparation."
- Praeger, A. H., "Notes on the Crystallization of High Grade Masecutes."
- Clayton, J. L., "An Aid to the Calculation of Pol in Juice Analysis."
- Clayton, J. L., "The Conditioning of Sugar for Bagging."

#### TECHNICAL COMMUNICATIONS.

In the current year the following Technical Communications were prepared and published by this Division:—

- 1939.—Technical Communication No. 5—"Subsidiary Performance Tests, 1938 Season," by G. H. Jenkins.
- Technical Communication No. 6—"Furnace Investigations, 1938 Season," by G. H. Jenkins.

Technical Communication No. 7—"Surplus Power from Surplus Bagasse," by G. H. Jenkins.

Technical Communication No. 8—"Masecuite Dilution Tests," by G. H. Jenkins.

Technical Communication No. 9—"Some Physical Properties of Molasses," by A. H. Praeger and J. L. Heron.

1940.—Technical Communication No. 1—"Cane Preparation. Part I.—The National Shredder," by E. R. Behne.

Technical Communication No. 2—"Rotary Filters for Treatment of Cane Muds," by G. H. Jenkins.

Technical Communication No. 3—"Automatic pH Control," by J. L. Clayton.

#### NEWS LETTER.

In order to provide a means of acquainting mill executives with the progress of experimental work carried out by the Technology Division and current literature not readily available to the mills, a "News Letter" service has been initiated. This News Letter, the first issue of which appeared in June, is to be forwarded to mills intermittently throughout the year, and is expected to provide a further means of contact between the Technology Division and the mills.

#### SLACK SEASON RESEARCH BY MILL OFFICERS.

Mr. J. Webster, of Racecourse Mill, and Mr. E. D. Jensen, of the Fairymead Sugar Company, carried out researches into the properties of raw sugar during the past slack season. Mr. Webster investigated the hygroscopic nature of raw sugar, and Mr. Jensen the determination of ash by conductometric methods. Technical Communications embodying the result of their works will be issued shortly.

#### SEASONAL INVESTIGATIONS, 1940.

The fourth annual meeting of the Mill Research Programme Committee was held on 8th April, 1940, during the Conference of the Queensland Society of Sugar Cane Technologists at Mackay. The new Chairman elected to succeed Mr. S. V. Fevre was Mr. M. A. Doolan, of Mulgrave. Mr. W. J. Ryan was elected Secretary.

The programme of investigation for the 1940 season adopted at this meeting is as follows:—

#### 1. Milling Tests.

The investigation into the preparation of cane to be extended to include:—

- (a) Further tests on the National Shredder;
- (b) The performance of Searby Shredders; and
- (c) Preparation with knives alone; comparative tests to be made with knives in different positions if practicable.

#### 2. Conditioning and Storage of Sugar.

Tests to be conducted in the Mackay District to investigate the possibility of conditioning sugar at the bagging station, to reduce the tendency towards subsequent caking.

Tests to be conducted into the factors which govern the deterioration of sugar in storage.

3. Cry

F  
cuits  
provid4. Filt  
Te

5. Mil

6. Ino  
AT  
only fT  
the p  
tons.  
than  
high  
recor  
ton of  
C  
juice  
Coeffi  
but b  
lowest  
C  
and c  
were  
and F  
at Me

19:

7

Tons o  
Tons o  
Pol in  
Fibre  
Purity  
P  
Cl  
S  
Gallon  
Appar  
Overal  
Recov  
BollinTons  
Tons  
Pol in  
Fibre  
Purity  
I  
S  
Gallo  
Appa  
Overl  
Reco  
Bollin

3. Crystallization Tests.

Following the work on crystallization of high-grade massecuites in 1939, the Webre pan to be tested as a means for providing rapid cooling of the massecuite prior to dropping.

4. Filter Tests.

- Tests to be carried out to determine: (a) The extent to which the operation of plate and frame filters may be improved by the use of filter-aids (especially bagacillo); (b) The optimum filtering cycle for such equipment.

5. Mill Visits.

6. Incidental Work, Unforeseen Work, and the following up of Advice Given.

COLORIMETRIC PH STANDARDS.

This year standard phenol red colour tubes were forwarded only to those mills which requested them.

MILL WORK—1939 SEASON.

The quantity of cane crushed in the 1939 season exceeded the previous record (established in 1938) by nearly 700,000 tons. The Southern and Central Districts crushed more cane than ever before, whilst in the North the crop was the third highest on record. The cane was of average quality and a record output of raw sugar resulted, the quantity of cane per ton of 94 n.t. sugar (6.77) having been bettered only in 1937.

Crushing rates again increased, whilst extraction and lost juice per cent. fibre both showed an improvement. The Coefficient of Work for 1939 was the highest recorded to date, but both the recovery and boiling-house efficiency were the lowest for a number of years.

Crushing for the 1939 season commenced on the 21st May and continued until 16th January. The first mills to start were Victoria and Macknade and the last to finish Pleystowe and Fairymead. The maximum harvesting period was 218 days at Macknade and the minimum 54 days at Eagleby.

Table with 5 columns: Season, Pol in Cane, Fibre in Cane, Purity, 1st Expd. Juice. Data is grouped by district: Southern District, Central District, Northern District, and All Districts, with rows for years 1930-1939.

TONS OF CANE PER TON OF 94 N.T. SUGAR.

Table with 12 columns for years 1928-1939, showing the amount of cane per ton of 94 n.t. sugar.

AVERAGE CRUSHING RATES (TONS CANE PER HOUR).

Table with 7 columns for years 1933-1939, showing average crushing rates in tons of cane per hour.

SOUTHERN DISTRICT.

Table with 8 columns for years 1933-1939, showing various metrics for the Southern District including tons of cane, sugar, fibre, and purity.

CENTRAL DISTRICT.

Table with 8 columns for years 1933-1939, showing various metrics for the Central District including tons of cane, sugar, fibre, and purity.

